

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

CLAIMS


We claim:

1. A compound of the structure



wherein R¹, R², R³, R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkoxy, cyano, nitro, cycloalkyl and carboxaldehyde;

with the proviso that at least one of R^1 or R^3 is



wherein D, B, Y and Z at each occurrence are independently selected from the group consisting of $-\text{CR}^6-$, $-\text{CR}^7\text{R}^8-$, $-\text{C}(\text{O})-$, $-\text{O}-$, $-\text{SO}_2-$, $-\text{S}-$, $-\text{N}=\text{}$, and $-\text{NR}^9-$;

n is an integer of zero to three;

R⁶, R⁷, R⁸ and R⁹, at each occurrence, are each independently selected from the group consisting of hydrogen, alkyl, carboxy,

hydroxyalkyl, alkylaminocarbonyl alkyl,

dialkylaminocarbonylalkyl and carboxyalkyl; and

R^{10} and R^{11} are each independently selected from the group consisting of

hydrogen, alkyl, cycloalkyl, alkoxyalkyl, alkoxycarbonylalkyl,

5 carboxyalkyl, hydroxyalkyl, heterocyclyl, heterocyclylalkyl and

heterocyclylamino;

wherein R^{10} and R^{11} may be joined to form a three to seven membered

heterocyclyl ring, said ring being optionally substituted with one or more
substituents R^{13} , wherein R^{13} , at each occurrence is independently selected

10 from the group consisting of alkyl, alkylene, alkoxy, alkoxyalkyl,
cycloalkyl, aryl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl,

heterocyclylalkylaminocarbonyl, hydroxy, hydroxyalkyl,

hydroxyalkoxyalkyl, carboxy, carboxyalkyl, carboxycarbonyl,

carboxaldehyde, alkoxycarbonyl, arylalkoxycarbonyl, aminoalkyl,

15 aminoalkanoyl, aminocarbonyl, carboxamido, alkoxycarbonylalkyl,

carboxamidoalkyl, cyano, tetrazolyl, alkanoyl, hydroxyalkanoyl,

alkanoyloxy, alkanoylamino, alkanoyloxyalkyl, alkanoylaminoalkyl,

sulfonate, alkylsulfonyl, alkylsulfonylaminocarbonyl,

arylsulfonylaminocarbonyl and heterocyclylsulfonylaminocarbonyl;

20 wherein A is an aryl or heterocyclyl group, said aryl or heterocyclyl group having at least

one substituent R^{12} , wherein R^{12} , at each occurrence, is independently selected


from the group consisting of hydrogen, halogen, alkyl, aryl, haloalkyl, hydroxy,

alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxyalkoxy, hydroxyalkyl, aminoalkyl,

aminocarbonyl, alkyl(alkoxycarbonylalkyl) aminoalkyl, heterocyclyl,
heterocyclylalkyl, carboxaldehyde, carboxaldehyde hydrazone, carboxamide,
alkoxycarbonylalkyl, carboxy, carboxyalkyl, carboxyalkoxy,
hydroxyalkylaminocarbonyl, cyano, amino, heterocyclylalkylamino,
5 carboxythioalkoxy, carboxycycloalkoxy, thioalkoxy, carboxyalkylamino, trans-
cinnamyl and heterocyclylalkylaminocarbonyl; and
wherein $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}$ and R^{13} are unsubstituted
or substituted with at least one electron donating or electron withdrawing
group;

10 or a pharmaceutically-acceptable salt, optical isomer or prodrug thereof.

2. The compound of claim 1 wherein R^3 is



D, B, Y and Z at each occurrence are independently selected from the
15 group consisting of $-CR^6=$, $-CR^7R^8-$, $-C(O)-$, $-O-$, $-SO_2-$, $-S-$,
 $-N=$, and $-NR^9-$;

n is an integer of zero to three;

R^6, R^7, R^8 and R^9 , at each occurrence, are each independently selected
from the group consisting of hydrogen, alkyl, carboxy,

hydroxyalkyl, alkylaminocarbonyl alkyl,
dialkylaminocarbonylalkyl and carboxyalkyl;

R^{10} and R^{11} are each independently selected from the group consisting of

hydrogen, alkyl, cycloalkyl, alkoxyalkyl, alkoxycarbonylalkyl,
5 carboxyalkyl, hydroxyalkyl, heterocyclyl, heterocyclylalkyl and
heterocyclylamino;

wherein R^{10} and R^{11} may be joined to form a three to seven membered

heterocyclyl ring, said ring optionally being substituted with one or more
substituents R^{13} , wherein R^{13} at each occurrence is independently selected

10 from the group consisting of alkyl, alkylene, alkoxy, alkoxyalkyl,
cycloalkyl, aryl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl,
heterocyclylalkylaminocarbonyl, hydroxy, hydroxyalkyl,
hydroxyalkoxyalkyl, carboxy, carboxyalkyl, carboxycarbonyl,
carboxaldehyde, alkoxycarbonyl, arylalkoxycarbonyl, aminoalkyl,
15 aminoalkanoyl, aminocarbonyl, carboxamido, alkoxycarbonylalkyl,
carboxamidoalkyl, cyano, tetrazolyl, alkanoyl, hydroxyalkanoyl,
alkanoyloxy, alkanoylamino, alkanoyloxyalkyl, alkanoylaminoalkyl,
sulfonate, alkylsulfonyl, alkylsulfonylaminocarbonyl,
arylsulfonylaminocarbonyl and heterocyclylsulfonylaminocarbonyl;

20 R^1 and R^2 are each independently selected from the group consisting of hydrogen,
halogen, haloalkyl and nitro; and

R^4 and R^5 are each independently selected from the group of hydrogen and alkyl.

wherein R^1 , R^2 , R^4 and R^5 are each independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkoxy, cyano, nitro, cycloalkyl and carboxaldehyde;

D, B, Y and Z at each occurrence are independently selected from the group consisting of $-CR^6=$, $-CR^7R^8-$, $-C(O)-$, $-O-$, $-SO_2-$, $-S-$, $-N=$, and $-NR^9-$.

n is an integer of zero to three;

wherein R^6 , R^7 , R^8 and R^9 , at each occurrence, are each independently selected from the group consisting of hydrogen, alkyl, carboxy, hydroxyalkyl, alkylaminocarbonyl alkyl, dialkylaminocarbonylalkyl and carboxyalkyl;

R^{10} and R^{11} are each independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkoxyalkyl, alkoxycarbonylalkyl, carboxyalkyl, hydroxyalkyl, heterocyclyl, heterocyclylalkyl and heterocyclylamino;

wherein R^{10} and R^{11} may be joined to form a three to seven membered heterocyclyl ring, said ring optionally being substituted with one or more

wherein R^1 , R^2 , R^4 and R^5 are each independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkoxy, cyano, nitro, cycloalkyl and carboxaldehyde;

D, B, Y and Z at each occurrence are independently selected from the group consisting of $-CR^6=$, $-CR^7R^8-$, $-C(O)-$, $-O-$, $-SO_2-$, $-S-$, $-N=$, and $-NR^9-$.

n is an integer of zero to three;

wherein R^6 , R^7 , R^8 and R^9 , at each occurrence, are each independently selected from the group consisting of hydrogen, alkyl, carboxy, hydroxyalkyl, alkylaminocarbonyl alkyl, dialkylaminocarbonylalkyl and carboxyalkyl;

R^{10} and R^{11} are each independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkoxyalkyl, alkoxycarbonylalkyl, carboxyalkyl, hydroxyalkyl, heterocyclyl, heterocyclylalkyl and heterocyclylamino;

wherein R^{10} and R^{11} may be joined to form a three to seven membered heterocyclyl ring, said ring optionally being substituted with one or more

substituents R^{13} , wherein R^{13} at each occurrence is independently selected from the group consisting of alkyl, alkylene, alkoxy, alkoxyalkyl, cycloalkyl, aryl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylalkylaminocarbonyl, hydroxy, hydroxyalkyl, hydroxyalkoxyalkyl, carboxy, carboxyalkyl, carboxycarbonyl, carboxaldehyde, alkoxycarbonyl, arylalkoxycarbonyl, aminoalkyl, aminoalkanoyl, aminocarbonyl, carboxamido, alkoxycarbonylalkyl, carboxamidoalkyl, cyano, tetrazolyl, alkanoyl, hydroxyalkanoyl, alkanoyloxy, alkanoylamino, alkanoyloxyalkyl, alkanoylaminoalkyl, sulfonate, alkylsulfonyl, alkylsulfonylaminocarbonyl, arylsulfonylaminocarbonyl and heterocyclylsulfonylaminocarbonyl;

R^{12} , at each occurrence, is independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkoxy, carboxyalkoxy, carboxyalkyl and heterocyclyl; and,

p is an integer of zero to five;

wherein R^1 , R^2 , R^4 , R^5 , R^{10} , R^{11} , R^{12} and R^{13} are unsubstituted or substituted with at least one electron donating group or electron withdrawing group.

4. The compound of claim 3 wherein p is one;

R^4 and R^5 are hydrogen;

R^{12} is selected from the group consisting of halogen, alkyl, alkoxy,

5 carboxyalkoxy, carboxyalkyl and heterocyclyl; and


R^{10} and R^{11} are joined to form a three to seven membered heterocyclyl ring; said

ring selected from the group consisting of piperidine, piperazine,

morpholine, pyrrolidine and azetidine.

10

5. The compound of claim 1 of the structure



wherein D and B are each independently selected from the group consisting of

$-N=$ and $-CR^6=$;

R^1 and R^2 are each independently selected from the group consisting of hydrogen,

15

halogen and haloalkyl;

R^{10} and R^{11} are each independently selected from the group consisting of

hydrogen, alkyl, cycloalkyl, alkoxyalkyl, alkoxycarbonylalkyl,

carboxyalkyl, hydroxyalkyl, heterocyclyl, heterocyclylalkyl and heterocyclylamino;

wherein R^{10} and R^{11} may be joined to form a three to seven membered

heterocyclyl ring, said ring optionally substituted with one or more
 5 substituents R^{13} , wherein R^{13} at each occurrence is independently selected from the group consisting of alkyl, alkylene, alkoxy, alkoxyalkyl, cycloalkyl, aryl, heterocyclyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylalkylaminocarbonyl, hydroxy, hydroxyalkyl, hydroxyalkoxyalkyl, carboxy, carboxyalkyl, carboxycarbonyl,
 10 carboxaldehyde, alkoxycarbonyl, arylalkoxycarbonyl, aminoalkyl, aminoalkanoyl, aminocarbonyl, carboxamido, alkoxycarbonylalkyl, carboxamidoalkyl, cyano, tetrazolyl, alkanoyl, hydroxyalkanoyl, alkanoyloxy, alkanoylamino, alkanoyloxyalkyl, alkanoylaminoalkyl, sulfonate, alkylsulfonyl, alkylsulfonylaminocarbonyl,
 15 arylsulfonylaminocarbonyl and heterocyclylsulfonylaminocarbonyl;

R^{12} , at each occurrence, is independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkoxy, carboxyalkoxy, carboxyalkyl and heterocyclyl; and,

p is an integer of zero to five;

20 wherein R^1 , R^2 , R^{10} , R^{11} , R^{12} and R^{13} are unsubstituted or substituted with at least one electron donating group or electron withdrawing group.

6. The compound of claim 5 wherein p is one;

5 R¹² is selected from the group consisting of halogen, alkyl, alkoxy,
carboxyalkoxy, carboxyalkyl and heterocyclyl; and

R¹⁰ and R¹¹ are joined to form a three to seven membered heterocyclyl ring; said
ring selected from the group consisting of piperidine, piperazine,
morpholine, pyrrolidine and azetidine.

10

7. The compound of claim 1 selected from the group consisting of 1-(6-(4-(2-
isopropyl-phenylsulfanyl)-3-trifluoromethyl-phenyl)-pyrimidin-4-yl)-piperidine-3-
carboxylic acid, 4-(4-(2-isopropyl-phenylsulfanyl)-3-trifluoromethyl-phenyl)-6-(3-(2*H*-
tetrazol-5-yl)-piperidin-1-yl)-pyrimidine, 4-(4-(2-isopropyl-phenylsulfanyl)-3-

15 trifluoromethyl-phenyl)-6-(4-(2*H*-tetrazol-5-yl)-piperidin-1-yl)-pyrimidine, (1-(6-(4-(2-
isopropyl-phenylsulfanyl)-3-trifluoromethyl-phenyl)-pyrimidin-4-yl)-piperidin-3-yl)-
methanol, 2-(1-(6-(4-(2-isopropyl-phenylsulfanyl)-3-trifluoromethyl-phenyl)-pyrimidin-4-
yl)-piperidin-4-yl)-ethanol, *N*-(1-(4-(4-(2-isopropyl-phenylsulfanyl)-3-trifluoromethyl-
phenyl)-pyridin-2-yl)-pyrrolidin-3-yl)-acetamide, 1-(4-(4-(2-methoxy-phenylsulfanyl)-3-
20 trifluoromethyl-phenyl)-pyridin-2-yl)-pyrrolidine-3-ol,

N-1-(4-(4-(2-methoxy-phenylsulfanyl)-3-trifluoromethyl-phenyl)-pyridin-2-yl)-
pyrrolidine-3-yl)-acetamide, *N*-1-(4-(4-(2-methoxy-phenylsulfanyl)-3-trifluoromethyl-
phenyl)-pyridin-2-yl)-pyrrolidine-3-yl)-acetamide, *N*-(1-(4-(4-(2,3-dihydro-

- benzo(1,4)dioxin-6-ylsulfanyl)-3-trifluoromethyl-phenyl)-pyridin-2-yl)-pyrrolidin-3-yl)-
acetamide, 4'-(4-(2,3-dihydro-benzo(1,4)dioxin-6-ylsulfanyl)-3-trifluoromethyl-phenyl)-
3,4,5,6-tetrahydro-2*H*-(1,2')bipyridinyl-4-carboxylic acid and 4'-(4-(2,3-dihydro-
benzo(1,4)dioxin-6-ylsulfanyl)-3-trifluoromethyl-phenyl)- 3,4,5,6-tetrahydro-2*H*-
5 (1,2')bipyridinyl-3-carboxylic acid.

8. A composition comprising:

a compound of claim 1

in a pharmaceutically acceptable carrier.

10

9. A method of inhibiting inflammation or suppressing immune response in a
mammal comprising administering to said mammal a therapeutic amount of a
compound of claim 1.

15